

Contents

Volume 106 (1) 1989

Patrick, T. D., Lewer, C. E. and Pain, V. M. Preparation and characterization of cell-free protein synthesis systems from oocytes and eggs of <i>Xenopus laevis</i>	1	Karr, T. L. and Kornberg, T. B. <i>fushi tarazu</i> protein expression in the cellular blastoderm of <i>Drosophila</i> detected using a novel imaging technique	95
Van den Eynde, H., Mazabraud, A. and Denis, H. Biochemical research on oogenesis. RNA accumulation in the oocytes of the newt <i>Pleurodeles waltl</i>	11	Bissen, S. T. and Weisblat, D. A. The duration and compositions of cell cycles in embryos of the leech, <i>Helobdella triserialis</i>	105
Garzino, V., Berenger, H. and Pradel, J. Expression of laminin and of a laminin-related antigen during early development of <i>Drosophila melanogaster</i>	17	Gard, A. L. and Pfeiffer, S. E. Oligodendrocyte progenitors isolated directly from developing telencephalon at a specific phenotypic stage: myelinogenic potential in a defined environment	119
Fundele, R., Norris, M. L., Barton, S. C., Reik, W. and Surani, M. A. Systematic elimination of parthenogenetic cells in mouse chimaeras	29	Maine, E. M. and Kimble, J. Identification of genes that interact with <i>glp-1</i> , a gene required for inductive cell interactions in <i>Caenorhabditis elegans</i>	133
Beddington, R. S. P., Morgernstern, J., Land, H. and Hogan, A. An <i>in situ</i> transgenic enzyme marker for the midgestation mouse embryo and the visualization of inner mass clones during early organogenesis	37	Ambrosio, L., Mahowald, A. P. and Perrimon, N. <i>l(1)pole hole</i> is required maternally for pattern formation in the terminal regions of the embryo	145
Knox, J. P., Day, S. and Roberts, K. A set of cell surface glycoproteins forms an early marker of cell position, but not cell type, in the root apical meristem of <i>Daucus carota</i> L.	47	Bloom, T. L. The effects of phorbol ester on mouse blastomeres: a role for protein kinase C in compaction	159
Simpson, P. and Carteret, C. A study of <i>shaggy</i> reveals spatial domains of expression of <i>achaete-scute</i> alleles on the thorax of <i>Drosophila</i>	57	Ruiz I Altaba, A. and Melton, D. A. Bimodal and graded expression of the <i>Xenopus</i> homeobox gene <i>Xhox3</i> during embryonic development	173
Taylor, M., Treisman, R., Garrett, N. and Mohun, T. Muscle-specific (CARG) and serum-responsive (SRE) promoter elements are functionally interchangeable in <i>Xenopus</i> embryos and mouse fibroblasts	67	Anderson, H. and Tucker, R. P. Spatial and temporal variation in the structure of the basal lamina in embryonic grasshopper limbs during pioneer neurone outgrowth	185
Paterno, G. D., Gillespie, L. L., Dixon, M. S., Slack, J. M. W. and Heath, J. K. Mesoderm-inducing properties of INT-2 and kFGF: two oncogene-encoded growth factors related to FCF	79	Sabbag, K. R., Casanova, J. E. and Gabel, L. B. Plasminogen activator expression in F9 teratocarcinoma embryoid bodies and their endoderm derivatives	195
Kalcheim, C. and Teillet, M.-A. Consequences of somite manipulation on the pattern of dorsal root ganglion development	85	Gillespie, L. L., Paterno, G. D. and Slack, J. M. W. Analysis of competence: receptors for fibroblast growth factor in early <i>Xenopus</i> embryos	203