

# Contents

Volume 110 (2) 1990

- Sullivan, W., Minden, J. S. and Alberts, B. M.**  
*daughterless-abo-like*, a *Drosophila* maternal-effect mutation that exhibits abnormal centrosome separation during the late blastoderm divisions 311
- Hemmati-Brivanlou, A., Frank, D., Bolce, M. E., Brown, R. D., Sive, H. L. and Harland, R. M.**  
Localization of specific mRNAs in *Xenopus* embryos by whole-mount *in situ* hybridization 325
- Ichijo, H., Fujita, S., Matsuno, T. and Nakamura, H.**  
Rotation of the tectal primordium reveals plasticity of target recognition in retinotectal projection 331
- Dencker, L., Annerwall, E., Busch, C. and Eriksson, U.**  
Localization of specific retinoid-binding sites and expression of cellular retinoic-acid-binding protein (CRABP) in the early mouse embryo 343
- Altmann, G. G. and Quaroni, A.**  
Behavior of fetal intestinal organ culture explanted onto a collagen substratum 353
- Vaessen, M.-J., Meijers, J. H. C., Bootsma, D. and van Kessel, A. G.**  
The cellular retinoic-acid-binding protein is expressed in tissues associated with retinoic-acid-induced malformations 371
- Ohneda, O., Yanai, N. and Obinata, M.**  
Microenvironment created by stromal cells is essential for a rapid expansion of erythroid cells in mouse fetal liver 379
- Renfree, M. B., Robinson, E. S., Short, R. V. and Vandenberg, J. L.**  
Mammary glands in male marsupials: I. Primordia in neonatal opossums *Didelphis virginiana* and *Monodelphis domestica* 385
- Tesarik, J., Drahorad, J., Testart, J. and Mendoza, C.**  
Acrosin activation follows its surface exposure and precedes membrane fusion in human sperm acrosome reaction 391
- Wehrle, B. and Chiquet, M.**  
Tenascin is accumulated along developing peripheral nerves and allows neurite outgrowth *in vitro* 401
- Charlebois, T. S., Henry, J. J. and Grainger, R. M.**  
Differential cytokeratin gene expression reveals early dorsal-ventral regionalization in chick mesoderm 417
- Ovsenek, N. and Heikkila, J. J.**  
DNA sequence-specific binding activity of the heat-shock transcription factor is heat-inducible before the midblastula transition of early *Xenopus* development 427
- Albano, R. M., Godsave, S. F., Huylebroeck, D., van Nimmen, K., Isaacs, H. V., Slack, J. M. W. and Smith, J. C.**  
A mesoderm-inducing factor produced by WEHI-3 murine myelomonocytic leukemia cells is activin A 435
- Gatherer, D., ten Dijke, P., Baird, D. T. and Akhurst, R. J.**  
Expression of TGF- $\beta$  isoforms during first trimester human embryogenesis 445
- Otte, A. P., Kramer, I. M., Mannesse, M., Lambrechts, C. and Durston, A. J.**  
Characterization of protein kinase C in early *Xenopus* embryogenesis 461
- Bennett, D. C., Huszar, D., Laipis, P. J., Jaenisch, R. and Jackson, I. J.**  
Phenotypic rescue of mutant brown melanocytes by a retrovirus carrying a wild-type tyrosinase-related protein gene 471
- Thompson, E. M., Fleming, K. A., Evans, D. J., Fundele, R., Surani, M. A. and Wright, N. A.**  
Gastric endocrine cells share a clonal origin with other gut cell lineages 477
- Kucera, J. and Walro, J. M.**  
Treatment with  $\beta$  bungarotoxin blocks muscle spindle formation in fetal rats 483
- Metcalfe, W. K., Myers, P. Z., Trevarrow, B., Bass, M. B. and Kimmel, C. B.**  
Primary neurons that express the L2/HNK-1 carbohydrate during early development in the zebrafish 491

Continued overleaf

|  |     |   |     |
|--|-----|---|-----|
| <b>Tuan, R. S. and Bigioni, N.</b><br>Ca <sup>2+</sup> -activated ATPase of the mouse<br>chorioallantoic placenta: developmental<br>expression, characterization and<br>cytohistochemical localization   | 505 | <b>Tanaka, H., Kinutani, M., Agata, A.,<br/>Takashima, Y. and Obata, K.</b><br>Pathfinding during spinal tract formation in the<br>chick-quail chimera analysed by species-specific<br>monoclonal antibodies                                | 565 |
| <b>Harper, S. and Davies, A. M.</b><br>NGF mRNA expression in developing cutaneous<br>epithelium related to innervation density  | 515 | <b>Huang, J. Q., Turbide, C., Daniels, E., Jothy, S.<br/>and Beauchemin, N.</b><br>Spatiotemporal expression of murine<br>carcinoembryonic antigen (CEA) gene family<br>members during mouse embryogenesis                                  | 573 |
| <b>Ginsburg, M., Snow, M. H. L. and McLaren, A.</b><br>Primordial germ cells in the mouse embryo<br>during gastrulation  | 521 | <b>Frohman, M. A., Boyle, M. and Martin, G. R.</b><br>Isolation of the mouse <i>Hox-2.9</i> gene; analysis of<br>embryonic expression suggests that positional<br>information along the anterior-posterior axis is<br>specified by mesoderm | 589 |
| <b>Nadon, N. L., Duncan, I. D. and Hudson, L. D.</b><br>A point mutation in the proteolipid protein gene<br>of the 'shaking pup' interrupts oligodendrocyte<br>development   | 529 | <b>Pelton, R. W., Dickinson, M. E., Moses, H. L.<br/>and Hogan, B. L. M.</b><br><i>In situ</i> hybridization analysis of TGFβ3 RNA<br>expression during mouse development:<br>comparative studies with TGFβ1 and β2                         | 609 |
| <b>Pullar, D., te Kronnie, G., Peiris, D.,<br/>Taverne, N., Jeacock, M. K., Stroband, H. W. J.<br/>and Shepherd, D. A. L.</b><br>Morphological and radiochemical evidence for the<br>metabolism of exogenous proteins by the<br>preimplantation sheep blastocyst | 539 | <b>Casanova, J.</b><br>Pattern formation under the control of the<br>terminal system in the <i>Drosophila</i> embryo  | 621 |
| <b>Saggin, L., Gorza, L., Ausoni, S.<br/>and Schiaffino, S.</b><br>Cardiac troponin T in developing, regenerating<br>and denervated rat skeletal muscle  | 547 | <b>Erselius, J. R., Goulding, M. D. and Gruss, P.</b><br>Structure and expression pattern of the murine<br><i>Hox-3.2</i> gene  | 629 |
| <b>Hahnel, A. C., Rappolee, D. A., Millan, J. L.,<br/>Manes, T., Ziomek, C. A., Theodosiou, N. G.,<br/>Werb, Z., Pedersen, R. A. and Schultz, G. A.</b><br>Two alkaline phosphatase genes are expressed<br>during early development in the mouse embryo          | 555 | <b>Plachov, D., Chowdhury, K., Walther, C.,<br/>Simon, D., Guenet, J.-L. and Gruss, P.</b><br><i>Pax8</i> , a murine paired box gene expressed in the<br>developing excretory system and thyroid gland                                      | 643 |